

Vegetation changes during the second part of the Wurmian glaciation in lowland and mountain regions: refugia and fire history

Thesis of Doctoral (PhD) dissertation

Ildikó Vincze



Eötvös Loránd University, Faculty of Science

Doctoral School of Environmental Sciences

Leader: Prof. Imre Jánosi

Thematic Program in Environmental Earth Sciences

Leader: Prof. Csaba Szabó

Supervisor: Enikő Magyari, DSc

Department of Environmental and Landscape Geography, Eötvös Loránd University

MTA-MTM-ELTE Research Group for Paleontology

2019

1 Introduction, aims

The motivation in the background of my doctoral research was the scarcity of plant macrofossil studies that would cover the Last Glacial Maximum and the Late Pleniglacial intervals (24000–14600 cal yr BP, TZEDAKIS et al. 2013) in the Carpathian Region. The available pollen and charcoal studies from Eastern and Southern Europe imply sparsely wooded landscape during this period (e.g. WILLIS et al. 2000; WILLIS & VAN ANDEL 2004; TANTAU et al. 2006; TONKOV et al. 2011; GAŁKA et al. 2013; TZEDAKIS et al. 2013). On the other hand, plant macrofossil studies from this period are scarce, therefore there is little evidence on the local presence of a species soon after the LGM that would confirm the refugial role of the Carpathians during the Last Glacial Maximum. Several studies investigated the changes of the vegetation and environment during the LPG in Hungary (e.g. BORSY 1989; CSERNY 2002; CSINÁDY 1960; JAKAB et al. 2005; JAKAB et al. 2009; MAGYARI et al. 1999; MAGYARI et al. 2000; MEDZIHRADSZKY & BAJZÁTH 1998; RUDNER & SÜMEGI 2001; SÜMEGI & GULYÁS 2004; SÜMEGI et al. 2013, WILLIS et al. 1995) that were mainly based on pollen and/or plant macrofossil analyses.

Pollen based quantitative climate reconstructions provide the regional aspect of the vegetation and climate change (SEPPÄ et al. 2004; WHITMORE et al. 2005). Due to their low taxonomic resolution these reconstructions can only carefully be applied in treeless environments (AARNES et al. 2012). On the other hand, plant macrofossils provide identification on species level, which would imply their local presence, as macrophytes are widely recognized as climate indicators due to their ability to react faster to climate change (IVERSEN 1954). Plant macrofossil studies are useful for paleoclimate and vegetation reconstructions because the environmental tolerance of living species that characterised the last glacial landscape are well known (VÄLIRANTA et al. 2015). The characteristic features of plant macrofossils helped in my doctoral research to investigate past local vegetation changes in lowlands and mountains, to reveal mire development phases, to detect local fire histories, to explore the effect of the Heinrich-1 cooling event on the vegetation (17500–16000 cal yr BP) and to reveal past treeline and timberline changes at high altitudes, species compositional changes in the treeline ecotone and the impact of human presence in the alpine region. The investigation of the Holocene treeline and timberline changes in the South Carpathians is highly significant due to the lack of these studies in the Romanian Carpathians, this topic

became more highlighted only recent past, when detailed study was made in the Rodna Mts, these results are also well comparable to the Alps, where treeline and timberline changes are extensively documented.

2 Study area and sampling

My doctoral research focused on three different study areas. I studied the deposits of a volcanic lake, Lake St. Anne (950 m asl.) in the Ciomadul Mts of the Eastern Carpathians, a small mire from the Érmellék region called Kokad (112 m asl.) and the high-altitude lakes of the Retezat Mts of the South Carpathians: Lake Brazi, Lake Gales, Lake Lia and Lake Bucura (from 1740 m to 2040 m a.s.l.). The sediment core of Kokad was taken in 2008, while the high-altitude lakes were sampled during 2007 and 2008. The coring of Lake St. Anne took place in 2010. The main focus of the selected cores was to cover the Late Pleniglacial and Lateglacial period by studying the vegetation changes during these intervals.

3 Methods

During the laboratory preparation of the samples, first heating with distilled water or 10% NaOH solution was applied, which was then followed by wet-sieving through 250µm and 180µm meshes. The plant macrofossil remains were studied under stereomicroscope with the aid of identification keys and the reference collection of the MTA-MTM-ELTE Research Group for Paleontology. Sediment components together with other remains (needles, seeds) were determined, as well as moss leaves and stems, monocotyledon remains, macrocharcoal were detected by using the methods of JAKAB et al. (2004). For the macrocharcoal analysis, 1cm³ samples were taken from the core continuously and prepared by following the methodology of GENRIES et al. (2012). Under the camera equipped binocular microscope not only the number of the macrocharcoal particles were detected, but their surface area too. The results were interpreted by the CharAnalysis software (HIGUERA et al. 2009). The plant macrofossil and macrocharcoal results were interpreted together with the results of the radiocarbon measurement, the loss-on-ignition, the geochemical elements, the grain-size distribution and the pollen results.

4 Theses

1. On the basis of the plant macrofossil analysis of Lake St Anne, I proved that coniferous species were the first to appear on the crater slope during the Late Pleniglacial warming, pine (*Pinus* sp.) around 16200 cal yr BP, then spruce (*Picea abies*) around 15900 cal yr BP. The early colonisation of the conifers coincided well with the pollen results, which revealed boreal forest expansion with the mixing of *Betula-Pinus-Larix-Picea* individuals.
2. I proved that the Younger Dryas cooling is presented by a short-term decrease in the organic content of Lake St Anne, in parallel with frequent spruce (*Picea abies*) and birch (*Betula* sp.) remains between 13000 and 12500 cal yr BP, suggesting that they became more frequent around the lake, while Poaceae and *Artemisia* became dominant regionally at the same time.
3. On the basis of the plant macrofossil analysis of Kokad, I proved that mesotrophic conditions occurred with brown mosses during the Late Pleniglacial, when reed (*Phragmites australis*) and bulrush (*Typha angustifolia* és *T. latifolia*) were present in the mosaic-like shallow water-level conditions. The appearance of bulrush implies warm, >12–15,7°C July mean temperatures soon after the LGM in the Érmellék region (VINCZE et al. 2019).
4. I proved that the timing of biogenic carbonate formation at Kokad mire is one of the earliest detected in the Great Hungarian Plain (GHP) at 16170 cal yr BP. The major period of the biogenic carbonate formation (from 15200 cal yr BP) preceded other similar events identified in the GHP substantially, by ~3000 years (VINCZE et al. 2019).
5. By reconstructing the local fire history at Kokad mire, I proved that local fire history significantly differs from the regional fire history: Kokad had low fire activity between 15000 and 4000 cal yr BP, either due to the reed-dominated vegetation around the mire during the late glacial, or due to the frequency of reed-swamp areas in the Érmellék region that was likely characterised by high groundwater table during the lateglacial and Early Holocene.
6. I demonstrated that both the treeline and the timberline reached their maximum elevation between 8000 és 3200 cal yr BP in the Retezat Mts, and their altitudinal fluctuation was 100–150 m (VINCZE et al. 2017). On the southern slope of the Retezat, Dwarf pine (*Pinus mugo*) dominated the vegetation at the level of 2040 m a.s.l. until the mid-Holocene, while Swiss stone pine (*P. cembra*) and spruce (*Picea abies*) developed a closed forest at 1910 m a.s.l. (VINCZE et al. 2017).
7. By comparing the local fire history of the Retezat Mts (southern and northern slopes), I have proved that the fire activity on the southern slope agrees well with the regionally

reconstructed Early Holocene high fire activity, although the human impact only related to moderate biomass burning until 1300 cal yr BP and became more intensive during the last 1000 years.

8. I have recognized by comparing the plant macrofossil and pollen results that human impact in the Retezat Mts. probably became significant during the Bronze Age, from 4200 cal yr BP, and it significantly changed the species composition and the extension of the treeline ecotone zone (VINCZE et al. 2017). On the southern slope, the alpine meadows dominated by *Nardus stricta* and *Festuca* spp. today developed under continuous grazing over the past 2600–2200 years (VINCZE et al. 2017).

5 References

- AARNES, I., KÜHL, N., & BIRKS, H. H. 2012: Quantitative climate reconstruction from late-glacial and early Holocene plant macrofossils in western Norway using the probability density function approach. – *Review of Palaeobotany and Palynology* **170**, 27–39.
- BORSY, Z. 1989: Az Alföld hordalékkúpjainak negyedidőszaki fejlődéstörténete. – *Földrajzi Értesítő* **XXXVIII./3–4**, 211–224.
- CSERNY, T. 2002: A balatoni negyedidőszaki üledékek kutatási eredményei. – *Földtani Közöny* **132**, 193–213.
- CSINÁDY, G. 1960: A kokadi láp palynológiai vizsgálata. – *Acta Univ. Debrecen* **6/2**, 239–251.
- GÁLKA, M., TOBOLSKI, K., ZAWISZA, E., & GOSLAR, T. 2013: Postglacial history of vegetation, human activity and lake-level changes at Jezioro Linówek in northeast Poland, based on multi-proxy data. – *Vegetation History and Archaeobotany* **23/2**, 123–152.
- GENRIES, A., FINSINGER, W., ASNONG, H., BERGERON, Y., CARCAILLET, C., GARNEAU, M., HÉLY, C., & ALI, A. A. 2012: Local versus regional processes: can soil characteristics overcome climate and fire regimes by modifying vegetation trajectories? – *Journal of Quaternary Science* **27/7**, 745–756.
- HIGUERA, P. E., BRUBAKER, L. B., ANDERSON, P. M., HU, F. S., & BROWN, T. A. 2009: Vegetation mediated the impacts of postglacial climate change on fire regimes in the south-central Brooks Range, Alaska. – *Ecological Monographs* **79/2**, 201–219.
- IVERSEN, J. 1954: The late-glacial flora of Denmark and its relation to climate and soil. – *Danmarks geologiske undersøgelse* **2/80**, 87–119.
- JAKAB, G., SÜMEGI, P., & MAGYARI, E. 2004: A new paleobotanical method for the description of Late Quaternary organic sediments (Mire-development pathways and paleoclimatic records from S Hungary). – *Acta Geologica Hungarica* **47/4**, 373–409.
- JAKAB, G., SÜMEGI, P., & SZÁNTÓ, Z. 2005: Késő-glaciális és holocén vízszintingadozások a Szigligeti-öbölben (Balaton) makrofosszília vizsgálatok alapján. (Late glacial and Holocene water level changes in the Szigliget Bay, Lake Balaton based on macrofossil investigations). – *Földtani Közöny* **135/3**, 405–431.
- JAKAB, G., MAJKUT, P., JUHÁSZ, I., GULYÁS, S., SÜMEGI, P., TÖRÖCSIK, T. 2009: Palaeoclimatic signals and anthropogenic disturbances from the peatbog at Nagybárcány (North Hungary). – *Hydrobiologia* **631/1**, 87–106.
- MAGYARI, E., JAKAB, G., RUDNER, E., & SÜMEGI, P. 1999: Palynological and plant macrofossil data on Late Pleistocene short-term climatic oscillations in NE-Hungary. – *Acta Palaeobotanica. Supplement* **2**, 491–502.
- MAGYARI, E. K., JAKAB, G., SÜMEGI, P., RUDNER, E., & MOLNÁR, M. 2000: Paleobotanikai vizsgálatok a keleméri Mohos-tavakon (Palaeobotanical investigations on Mohos-lakes, Kelemér, N-Hungary). – *Tőzegmohás élőhelyek Magyarországon: kutatás, kezelés, védelem* 101–131.
- MEDZIHRADSKY, Z., & BAJZÁTH, J. 1998: The occurrence of arctic-alpine *Betula* species in the Hungarian Pleistocene. – *Annals His. Nat. Mus. Nat. Hung.* **90**, 27–33.
- RUDNER, Z. E., & SÜMEGI, P. 2001: Recurring Taiga forest-steppe habitats in the Carpathian Basin in the Upper Weichselian. – *Quaternary International* **76**, 177–189.

- SEPPÄ, H., BIRKS, H. J. B., ODLAND, A., POSKA, A., & VESKI, S. 2004: A modern pollen–climate calibration set from northern Europe: developing and testing a tool for palaeoclimatological reconstructions. – *Journal of Biogeography* **31/2**, 251–267.
- SÜMEGI, P., & GULYÁS, S. 2004: The geohistory of Bátorliget marshland: an example for the reconstruction of late quaternary environmental changes and past human impact from the northeastern part of the Carpathian Basin– Archaeolingua.
- SÜMEGI, P., MAGYARI, E., DÁNIEL, P., MOLNÁR, M., & TÖRÖCSIK, T. 2013: Responses of terrestrial ecosystems to Dansgaard–Oeshger cycles and Heinrich-events: A 28,000-year record of environmental changes from SE Hungary. – *Quaternary International* **293**, 34–50.
- TANTAU, I., REILLE, M., DE BEAULIEU, J.-L., & FARCAS, S. 2006: Late Glacial and Holocene vegetation history in the southern part of Transylvania (Romania): pollen analysis of two sequences from Avrig. – *Journal of Quaternary Science* **21/1**, 49–61.
- TONKOV, S., POSSNERT, G., & BOZILOVA, E. 2011: The Lateglacial in the Rila Mountains (Bulgaria) revisited: The pollen record of Lake Ribno (2184 m). – *Review of Palaeobotany and Palynology* **166/1–2**, 1–11.
- TZEDAKIS, P. C., EMERSON, B. C., & HEWITT, G. M. 2013: Cryptic or mystic? Glacial tree refugia in northern Europe. – *Trends in Ecology and Evolution* **28/12**, 696–704.
- VÄLIRANTA, M., SALONEN, J. S., HEIKKILÄ, M., AMON, L., HELMENS, K., KLIMASCHEWSKI, A., KUHY, P., KULTTI, S., POSKA, A., SHALA, S., VESKI, S., & BIRKS, H. H. 2015: Plant macrofossil evidence for an early onset of the Holocene summer thermal maximum in northernmost Europe. – *Nature Communications* **6**.
- VINCZE, I., FINSINGER, W., JAKAB, G., BRAUN, M., HUBAY, K., VERES, D., DELI, T., SZALAI, Z., SZABÓ, Z., & MAGYARI, E. 2019: Paleoclimate reconstruction and mire development in the Eastern Great Hungarian Plain for the last 20,000 years. – *Review of Palaeobotany and Palynology* 104112.
- VINCZE, I., ORBÁN, I., BIRKS, H. H., PÁL, I., FINSINGER, W., HUBAY, K., MARINOVA, E., JAKAB, G., BRAUN, M., BIRÓ, T., TÓTH, M., DANAU, C., FERENCZ, I. V., & MAGYARI, E. K. 2017: Holocene treeline and timberline changes in the South Carpathians (Romania): climatic and anthropogenic drivers on the southern slopes of the Retezat Mountains. – *The Holocene* **27** (11), 1613–1630.
- WHITMORE, J., GAJEWSKI, K., SAWADA, M., WILLIAMS, J. W., SHUMAN, B., BARTLEIN, P. J., MINCKLEY, T., VIAU, A. E., III, T. W., SHAFER, S., ANDERSON, P., & BRUBAKER, L. 2005: Modern pollen data from North America and Greenland for multi-scale paleoenvironmental applications. – *Quaternary Science Reviews* **24/16–17**, 1828–1848.
- WILLIS, K. J., SÜMEGI, P., BRAUN, M., & TÓTH, A. 1995: The late Quaternary environmental history of Bátorliget, N.E. Hungary. – *Palaeogeography, Palaeoclimatology, Palaeoecology* **118/1–2**, 25–47.
- WILLIS, K. J., RUDNER, E., & SÜMEGI, P. 2000: The full-glacial forests of central and southeastern Europe. – *Quaternary Research* **53/2**, 203–213.
- WILLIS, K. J., & VAN ANDEL, T. H. 2004: Trees or no trees? The environments of central and eastern Europe during the Last Glaciation. – *Quaternary Science Reviews* **23/23–24**, 2369–2387.

Publications in the topic of the doctoral study

5.1. Publications in the topic of the doctoral study as a corresponding author

- VINCZE, I., FINSINGER, W., JAKAB, G., BRAUN, M., HUBAY, K., VERES, D., DELI, T., SZALAI, Z., SZABÓ, Z., & MAGYARI, E. (2019). Paleoclimate reconstruction and mire development in the Eastern Great Hungarian Plain for the last 20,000 years. *Review of Palaeobotany and Palynology*, <https://doi.org/10.1016/j.revpalbo.2019.104112>.
- VINCZE, I., ORBÁN, I., BIRKS, H. H., PÁL, I., FINSINGER, W., HUBAY, K., MARINOVA, E., JAKAB, G., BRAUN, M., BIRÓ, T., TÓTH, M., DANAU, C., FERENCZ, I. V., MAGYARI, E. K. (2017). Holocene treeline and timberline changes in the South Carpathians (Romania): Climatic and anthropogenic drivers on the southern slopes of the Retezat Mountains. *The Holocene*, **27** (11), 1613–1630.

5.2. Publications as a co-author

- FEURDEAN, A., VANNIÈRE, B., FINSINGER, W., WARREN, D., CONNOR, S.C., FORREST, M., LIAKKA, J., PANAIT, A., WERNER, C., ANDRIČ, M., BOBEK, P., CARTER, V.A., DAVIS, B., DIACONU, A.-C., DIETZE, E., FEESER, I., FLORESCU, G., GAŁKA, M., GIESECKE, T., JAHNS, S., JAMRICHOVÁ, E., KAJUKALO, K., KAPLAN, J., KARPIŃSKA-KOŁACZEK, M., KOŁACZEK, P., KUNEŠ, P., KUPRIYANOV, D., LAMENTOWICZ, M., LEMMEN, C., MAGYARI, E.K., MARCISZ, K., MARINOVA, E., NIAMIR, A.,

- NOVENKO, E., OBREMSKA, M., PĖDZISZEWSKA, A., PFEIFFER, M., POSKA, A., RÖSCH, M., SŁOWIŃSKI, M., STANČIKAITĖ, M., SZAL, M., ŚWIĘTA-MUSZNICKA, J., TANȚAU, I., THEUERKAUF, M., TONKOV, S., VALKÓ, O., VASSILJEV, J., VESKI, S., **VINCZE, I.**, WACNIK, A., WIETHOLD, J., HICKLER, T., 2019. Fire risk modulation by long-term dynamics in land cover and dominant forest type in Eastern and Central Europe. *Biogeosciences Discuss.* 2019, 1–36. <https://doi.org/10.5194/bg-2019-260>
- MAGYARI, E.K., PÁL, I., **VINCZE, I.**, VERES, D., JAKAB, G., BRAUN, M., SZALAI, Z., SZABÓ, Z., KORPONAI, J. (2019) Warm Younger Dryas summers and early Late Glacial spread of temperate deciduous trees in the Pannonian Basin during the last glacial termination (20-9 kyr cal BP). *Quaternary Science Reviews.* <https://doi.org/10.1016/j.quascirev.2019.105980>
- MAGYARI, E., **VINCZE, I.**, ORBÁN, I., BÍRÓ, T., & PÁL, I. (2018). Timing of major forest compositional changes and tree expansions in the Retezat Mts during the last 16,000 years. *Quaternary International*, 477, 40-58.
- LENDVAY, B., BÁLINT, M., PÁL, I., **VINCZE, I.**, ORBÁN, I., & MAGYARI, E. K. (2018). Plant macrofossils from lake sediment as the material to assess ancient genetic diversity: Did deforestation influence Norway spruce (*Picea abies*) in the South Carpathians? *Quaternary International*, 477, 106-116.
- ORBÁN, I., BIRKS H. H., **VINCZE, I.**, FINSINGER W., PÁL I., MARINOVA, E., JAKAB G., BRAUN M., HUBAY K., BÍRÓ T., MAGYARI E. K. (2018): Treeline and timberline dynamics on the northern and southern slopes of the Retezat Mountains (Romania) during the late glacial and the Holocene. *Quaternary International*. DOI:10.1016/j.quaint.2017.03.012.
- FINSINGER W., FEVRE J., ORBÁN I., PÁL I., **VINCZE I.**, HUBAY K., BIRKS HH, BRAUN M., TÓTH M., MAGYARI EK (2018): Holocene fire-regime changes near the treeline in the Retezat Mts. (Southern Carpathians, Romania). *Quaternary International*. DOI: 10.1016/j.quaint.2016.04.029.
- PÁL, I., BUCZKÓ, K., **VINCZE, I.**, FINSINGER, W., BRAUN, M., BÍRÓ, T., MAGYARI, E. K. (2016): Terrestrial and aquatic ecosystem responses to early Holocene rapid climate change (RCC) events in the South Carpathian Mountains, Romania. *Quaternary International* DOI: 10.1016/j.quaint.2016.11.015
- PÁL, I., MAGYARI, E. K., BRAUN, M., **VINCZE, I.**, PÁLFY, J., MOLNÁR, M., FINSINGER, W. AND BUCZKÓ, K. (2016): Small-scale moisture availability increase during the 8.2 ka climatic event inferred from biotic proxy records in the South Carpathians (SE Romania). *The Holocene*. DOI: 10.1177/0959683616640039.
- MAGYARI, E. K., DEMÉNY, A., BUCZKÓ, K., KERN, Z., VENNEMANN, T., FÓRIZS, I., **VINCZE, I.**, BRAUN, M., KOVÁCS, J. I., UDVARDI, B., VERES, D., 2013: A 13,600-year diatom oxygen isotope record from the South Carpathians (Romania): Reflection of winter conditions and possible links with North Atlantic circulation changes, *Quaternary International*, 293, pp. 136-149.

5.3. Oral and poster presentations

- VINCZE I.**, MAGYARI E., BRAUN M., HUBAY, K., FINSINGER W., JAKAB G., SZALAI Z. (2018): öskörnyezeti rekonstrukció a Keleti-Kárpátok Szent Anna-tavának vizsgálata alapján. Program, Előadáskivonatok, Kirándulásvezető, **21. Magyar Őslénytani Vándorgyűlés**, 2018. május 24-26. Félixfürdő, pp. 32-33.
- VINCZE I.**, MAGYARI E., PÁL I., FINSINGER W. (2017): Erdőtűz rekonstrukciós vizsgálatok módszertana és a Kárpát-medence tüzeseményei tavi és lápi szelvények alapján. Program, Előadáskivonatok, Kirándulásvezető, **20. Magyar Őslénytani Vándorgyűlés**, 2017. május 25-27. Tata-Tardos, pp. 47-48.
- VINCZE I.**, MAGYARI E., JAKAB G., BRAUN M., SZALAI Z., FINSINGER W. (2016): Paleoökológiai rekonstrukció növényi makrofosszília és makropernye vizsgálatok alapján. Program, Előadáskivonatok, Kirándulásvezető, **19. Magyar Őslénytani Vándorgyűlés**, 2015. május 26-28. Kozárd, pp. 41-42.
- VINCZE I.**, MAGYARI E. (2016) A növénytakaró változásai a neolitikum idején Polgár térségében paleoökológiai

- vizsgálatok alapján, valamint ízelítő a legújabb paleolitikus faszénvizsgálatok eredményeiből. „**Földünk múltjai**” – **Az ember és a táj kapcsolata a történelmi időkben** – konferencia: előadás szekció, KE Földrajz és Környezettudományi Intézet (Eger).
- VINCZE I., MAGYARI E., JAKAB G., BRAUN M.** (2015): A Kokadi-láp multi-proxy vizsgálata – Környezet- és klímarekonstrukció növényi makrofosszília vizsgálat alapján Program, Előadáskivonatok, Kirándulásvezető, **18. Magyar Őslénytani Vándorgyűlés**, 2015. Május 14-16. Varbó-Fónagyság, pp. 37.
- VINCZE I., ORBÁN I., H. H. BIRKS, PÁL I., HUBAY K., E. MARINOVA, JAKAB G., BRAUN M., BIRÓ T., C. DĂNĂU, W. FINSINGER, MAGYARI E.** (2015): Késő-glaciális és holocén erdőhatár-változások a Déli- Kárpátokban: klimatikus és emberi hatás a Retezat-hegység déli lejtőjén. **Tájhasználat és tájvédelem – kihívások és lehetőségek.** A Budapesten 2015. május 21-23. között megrendezett VI. Magyar Tájökológiai Konferencia előadásainak kivonatai. pp.110.
- VINCZE I., ORBÁN I., MARINOVA E., JAKAB G., BIRKS H. H., FINSINGER W., MAGYARI E. K.** (2014): Drivers of the Holocene treeline and timberline changes in the Retezat Mountains (South Carpathians, Romania). Abstract Book. **2nd International Workshop – Late Pleistocene and Holocene climatic variability in the Carpathian-Balkan region (CBW2014)**, 6th – 9th November 2014, Cluj-Napoca, Romania, p. 183-184. DOI: <http://dx.doi.org/10.4316/GEOREVIEW.2014.0.0.246>
- VINCZE I., ORBÁN I., MARINOVA E., JAKAB G., BIRKS H. H., FINSINGER W., MAGYARI E. K.** (2014): Late-glacial and Holocene treeline changes in the Southern Carpathians (Romania): climatic and anthropogenic drivers of the vegetation changes. Abstract Book. **9th European Palaeobotany and Palynology Conference**, 26th – 31th August 2014, Padova, Italy, p. 299-300.
- VINCZE I., ORBÁN I., BIRKS, H., MARINOVA, E., JAKAB G., MAGYARI E.** (2014):Holocén erdőhatár változások a Déli-Kárpátok Retezat hegységében: különbségek a déli és az északi lejtő között. (Holocene treeline changes in the South Carpathians (Retezat Mts.): differences between north and south slopes. Program, Előadáskivonatok, Kirándulásvezető, **17. Magyar Őslénytani Vándorgyűlés**, 2013. május 29-31. Győr, pp. 37-38.
- ORBÁN I., VINCZE I., BIRKS, H., MARINOVA E., JAKAB G., MAGYARI E.**: Késő-glaciális és holocén erdőhatár változások a Retezat-hegységben (Late glacial and Holocene treeline changes in the Retezat Mountains). In: **MAGYARI E. (szerk.): Késő negyedkori kvantitatív környezeti és klímarekonstrukciók a Déli-Kárpátokban (Late Quaternary Qualitative environmental and climate reconstructions in the South Carpathians) PROLONG PROJEKT** előadói absztraktkötete. p. 14-15.
- FINSINGER W., VINCZE I., FEVRE J., MAGYARI E.**: Lokális és regionális erdőtűzek a Retezat-hegységben az elmúlt 16 000 évben (Local and regional fire histories in the Retezat Mountains during the last 16 000 years). In: **MAGYARI, E. (szerk.): Késő negyedkori kvantitatív környezeti és klímarekonstrukciók a Déli-Kárpátokban (Late Quaternary Qualitative environmental and climate reconstructions in the South Carpathians) PROLONG PROJEKT** előadói absztraktkötete. p. 23.
- VINCZE I., PÁL I., MAGYARI E.**: Üledék hozzáférhetőségi adatbázis a Retezat hegység 4 tavi üledékszelvényére vonatkozóan. (Sediment availability database for the 4 studied lakes in the Retezat Mts.) In: **MAGYARI E. (szerk.): Késő negyedkori kvantitatív környezeti és klímarekonstrukciók a Déli-Kárpátokban (Late Quaternary Qualitative environmental and climate reconstructions in the South Carpathians) PROLONG PROJEKT** előadói absztraktkötete. p. 24.
- VINCZE I., ORBÁN I., BIRKS H., MARINOVA E., JAKAB G., MAGYARI E.** (2014): Holocén erdőhatár változások a Déli-Kárpátokban: klimatikus és antropogén hatások a mai erdőhatár és fahatár összetételében.(Holocene treeline shifts in the South Carpathians: climatic and anthropogenic impacts on the treeline vegetation). In: **SCHMIDT D.– KOVÁCS M. – BARTHA D. (szerk.): X. Aktuális Flóra- és Vegetációkutatás a Kárpát-medencében nemzetközi konferencia** absztraktkötete. – Nyugat-magyarországi Egyetem Kiadó, Sopron, p. 41.
- VINCZE I., MAGYARI E., ORBÁN I., MARINOVA, E., BIRKS, H., PÁL I., FINSINGER W.** (2013) Holocén erdőhatár változás a Déli- Kárpátokban: multi-proxy eredmények gleccsertavi üledékek vizsgálata alapján. Program, Előadáskivonatok, Kirándulásvezető, **16. Magyar Őslénytani Vándorgyűlés**, 2013. május 23-25 Orfű, pp. 42-43.
- MAGYARI, E.K., BUCZKÓ, K., DEMÉNY, A., KERN, Z., VENNEMANN, T., FÓRIZS, I., VINCZE, I.** (2012) A 13,600-year diatom oxygen isotope record from the South Carpathians (Romania): reflection of winter conditions and possible links with North Atlantic circulation changes. **International Paleolimnology Symposium IPS2012** Glasgow SECC 21-24 August, 2012. Abstract Book, p. 22.
- MAGYARI, E., DEMÉNY, A., BUCZKÓ, K., KERN, Z., VENNEMANN, T., FÓRIZS, I., VINCZE, I., BRAUN, M., KOVÁCS, J.I., VERES, D.** (2012) 13600 éves diatóma alapú oxigén izotóp adatsor a Déli-Kárpátokból: a téli félév éghajlatának rekonstrukciója és kapcsolata az észak-atlanti oszcillációval. **PAGES 2012 előadói absztraktkötet**, MTA CSFK Földtani és Geokémiai Intézet, 2012. április 11.
- MAGYARI, E. K., DEMÉNY, A., BUCZKÓ, K., KERN, Z., VENNEMANN, T., FÓRIZS, I., VINCZE, I., BRAUN, M., KOVÁCS,**

- J.I., VERES, D. (2012) 13600 éves diatóma alapú oxigén izotóp adatsor a Déli-Kárpátokból: a téli félév éghajlatának rekonstrukciója és kapcsolata az észak-atlanti oszcillációval. **15. Magyar Őslénytani Vándorgyűlés.** Program, Előadáskivonatok, Kirándulásvezető, p. 17.
- PÁL, I., VINCZE, I., FERENCZY GY., BRAUN, M., MOLNÁR, M., MAGYARI, E. (2012) Rövid távú holocén klímafluktuációk vegetációra gyakorolt hatása a Déli-Kárpátok Retyezát hegységében. **15. Magyar Őslénytani Vándorgyűlés.** Program, Előadáskivonatok, Kirándulásvezető, p. 20-21.
- VINCZE, I., PÁL, I., VENNEMANN, T., DEMÉNY, A., BUCZKÓ, K., FÓRIZS, I., KERN, Z., BRAUN, M., MAGYARI, E. (2012) Késő - glaciális és holocén oxigénizotóp-alapú klímarekonstrukció hibahatár becslése a Déli-Kárpátokban tavi üledékek elemzése alapján. **15. Magyar Őslénytani Vándorgyűlés.** Program, Előadáskivonatok, Kirándulásvezető, p. 28- 29.